

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number
WO 02/096905 A1

(51) International Patent Classification⁷: **C07D 417/04**,
A61K 31/427, A61P 03/10

(21) International Application Number: PCT/US02/16352

(22) International Filing Date: 23 May 2002 (23.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/295,196 1 June 2001 (01.06.2001) US

(71) Applicant (for all designated States except US): **VERTEX PHARMACEUTICALS INCORPORATED** [US/US];
Patent Department, 130 Waverly Street, Cambridge, MA 02139-4242 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **COCHRAN, John** [US/US]; 4 Philips Farm Road, Marshfield, MA 02050 (US). **NANTHAKUMAR, Suganthini** [LK/US]; 253 Spiers Road, Newton, MA 02459 (US). **HARRINGTON, Edmund** [IE/US]; 460 East 8th Street, South Boston, MA 02127 (US). **WANG, Jian** [CN/US]; 25 South Point Drive, Boston, MA 02125 (US).

(74) Agents: **ROBIDOUX, Andrea** et al.; Vertex Pharmaceuticals Inc., 130 Waverly Street, Cambridge, MA 02139-4242 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: THIAZOLE COMPOUNDS USEFUL AS INHIBITORS OF PROTEIN KINASES

(57) Abstract: The present invention provides a compound of formula I: or a pharmaceutically acceptable derivative thereof. These compounds are inhibitors of protein kinases, particularly inhibitors of GSK3, Aurora2, and Syk mammalian protein kinases. The invention also provides pharmaceutically acceptable compositions comprising the compounds of the invention and methods of utilizing those compounds and compositions in the treatment of various protein kinase mediated disorders.

WO 02/096905 A1

THIAZOLE COMPOUNDS USEFUL AS INHIBITORS OF PROTEIN KINASECROSS REFERENCE TO RELATED APPLICATION

This application claims priority to US Provisional Patent Application 60/295,158 filed June 1, 2001, the contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to compounds that are protein kinase inhibitors, pharmaceutically acceptable compositions comprising said compounds, and methods of use thereof. More particularly, the compounds are inhibitors of GSK-3, Aurora2, and Syk protein kinases and are useful for treating, or lessening the severity of, a variety of diseases and conditions, such as diabetes, Alzheimer's disease, stroke, proliferative disorders, and asthma.

10

BACKGROUND OF THE INVENTION

The search for new therapeutic agents has been greatly aided in recent years by better understanding of the structure of enzymes and other biomolecules associated with target diseases. One important class of enzymes that has been the subject of extensive study is the protein kinases.

Protein kinases mediate intracellular signal transduction. They do this by affecting a phosphoryl transfer from a nucleoside triphosphate to a protein acceptor that is involved in a signaling pathway. There

20

-2-

are a number of kinases and pathways through which extracellular and other stimuli cause a variety of cellular responses to occur inside the cell. Examples of such stimuli include environmental and chemical stress
5 signals (e.g. osmotic shock, heat shock, ultraviolet radiation, bacterial endotoxin, H_2O_2), cytokines (e.g. interleukin-1 (IL-1) and tumor necrosis factor α (TNF- α)), and growth factors (e.g. granulocyte macrophage-colony-stimulating factor (GM-CSF), and fibroblast growth
10 factor (FGF)). An extracellular stimulus may effect one or more cellular responses related to cell growth, migration, differentiation, secretion of hormones, activation of transcription factors, muscle contraction, glucose metabolism, control of protein synthesis and
15 regulation of cell cycle.

Many diseases and conditions are associated with abnormal cellular responses triggered by protein kinase-mediated events. These diseases include autoimmune diseases, inflammatory diseases, neurological
20 and neurodegenerative diseases, cancer, cardiovascular diseases, allergies and asthma, Alzheimer's disease or hormone-related diseases. Accordingly, there has been a substantial effort in medicinal chemistry to find protein kinase inhibitors that are effective as therapeutic
25 agents.

Glycogen synthase kinase-3 (GSK-3) is a serine/threonine protein kinase comprised of α and β isoforms that are each encoded by distinct genes [Coghlan et al., *Chemistry & Biology*, 7, 793-803 (2000); Kim and
30 Kimmel, *Curr. Opinion Genetics Dev.*, 10, 508-514 (2000)]. GSK-3 has been implicated in various diseases including diabetes, Alzheimer's disease, CNS disorders such as

-3-

manic depressive disorder and neurodegenerative diseases, and cardiomyocyte hypertrophy [WO 99/65897; WO 00/38675; and Haq et al., *J. Cell Biol.* (2000) 151, 117]. These diseases may be caused by, or result in, the abnormal operation of certain cell signaling pathways in which GSK-3 plays a role. GSK-3 has been found to phosphorylate and modulate the activity of a number of regulatory proteins. These include glycogen synthase which is the rate limiting enzyme necessary for glycogen synthesis, the microtubule associated protein Tau, the gene transcription factor β -catenin, the translation initiation factor eIF2B, as well as ATP citrate lyase, axin, heat shock factor-1, c-Jun, c-Myc, c-Myb, CREB, and CEPB α . These diverse targets implicate GSK-3 in many aspects of cellular metabolism, proliferation, differentiation and development.

In a GSK-3 mediated pathway that is relevant for the treatment of type II diabetes, insulin-induced signaling leads to cellular glucose uptake and glycogen synthesis. Along this pathway, GSK-3 is a negative regulator of the insulin-induced signal. Normally, the presence of insulin causes inhibition of GSK-3 mediated phosphorylation and deactivation of glycogen synthase. The inhibition of GSK-3 leads to increased glycogen synthesis and glucose uptake [Klein et al., *PNAS*, 93, 8455-9 (1996); Cross et al., *Biochem. J.*, 303, 21-26 (1994); Cohen, *Biochem. Soc. Trans.*, 21, 555-567 (1993); Massillon et al., *Biochem J.* 299, 123-128 (1994)]. However, in a diabetic patient where the insulin response is impaired, glycogen synthesis and glucose uptake fail to increase despite the presence of relatively high blood levels of insulin. This leads to abnormally high blood levels of glucose with acute and long term effects that

-4-

may ultimately result in cardiovascular disease, renal failure and blindness. In such patients, the normal insulin-induced inhibition of GSK-3 fails to occur. It has also been reported that in patients with type II
5 diabetes, GSK-3 is overexpressed [WO 00/38675]. Therapeutic inhibitors of GSK-3 are therefore potentially useful for treating diabetic patients suffering from an impaired response to insulin.

GSK-3 activity has also been associated with
10 Alzheimer's disease. This disease is characterized by the well-known β -amyloid peptide and the formation of intracellular neurofibrillary tangles. The neurofibrillary tangles contain hyperphosphorylated Tau protein where Tau is phosphorylated on abnormal sites.
15 GSK-3 has been shown to phosphorylate these abnormal sites in cell and animal models. Furthermore, inhibition of GSK-3 has been shown to prevent hyperphosphorylation of Tau in cells [Lovestone et al., *Current Biology* 4, 1077-86 (1994); Brownlee et al., *Neuroreport* 8, 3251-55
20 (1997)]. Therefore, it is believed that GSK-3 activity may promote generation of the neurofibrillary tangles and the progression of Alzheimer's disease.

Another substrate of GSK-3 is β -catenin which is degraded after phosphorylation by GSK-3. Reduced
25 levels of β -catenin have been reported in schizophrenic patients and have also been associated with other diseases related to increase in neuronal cell death [Zhong et al., *Nature*, 395, 698-702 (1998); Takashima et al., *PNAS*, 90, 7789-93 (1993); Pei et al., *J. Neuropathol. Exp*, 56, 70-78 (1997)].
30

Aurora-2 is a serine/threonine protein kinase that has been implicated in human cancer, such as colon,

-5-

breast and other solid tumors. This kinase is involved in protein phosphorylation events that regulate the cell cycle. Specifically, Aurora-2 plays a role in controlling the accurate segregation of chromosomes

5 during mitosis. Misregulation of the cell cycle can lead to cellular proliferation and other abnormalities. In human colon cancer tissue, the aurora-2 protein has been found to be overexpressed [Bischoff et al., *EMBO J.*, 17, 3052-3065 (1998); Schumacher et al., *J. Cell Biol.*, 143, 1635-1646 (1998); Kimura et al., *J. Biol. Chem.*, 272, 13766-13771 (1997)].

10

Syk is a tyrosine kinase that plays a critical role in FcεRI mediated mast cell degranulation and eosinophil activation. Accordingly, Syk kinase is

15 implicated in various allergic disorders, in particular asthma.

It has been shown that Syk binds to the phosphorylated gamma chain of the FcεRI receptor via N-terminal SH2 domains and is essential for downstream

20 signaling [Taylor et al, *Mol Cell Biol* 1995; 15:4149].

Inhibition of eosinophil apoptosis has been proposed as key mechanisms for the development of blood and tissue eosinophilia in asthma. IL-5 and GM-CSF are upregulated in asthma and are proposed to cause blood and

25 tissue eosinophilia by inhibition of eosinophil apoptosis. Inhibition of eosinophil apoptosis has been proposed as a key mechanism for the development of blood and tissue eosinophilia in asthma. It has been reported that Syk kinase is required for the prevention of

30 eosinophil apoptosis by cytokines (using antisense) [Yousefi et al, *J Exp Med* 1996;183:1407].


-6-

The role of Syk in FcγR dependent and independent response in bone marrow derived macrophages has been determined by using irradiated mouse chimeras reconstituted with fetal liver cells from Syk -/- embryos. Syk deficient macrophages were defective in phagocytosis induced by FcγR but showed normal phagocytosis in response to complement [Kiefer et al, Mol Cell Biol 1998; 18:4209]. It has also been reported that aerosolized Syk antisense suppresses Syk expression and mediator release from macrophages [Stenton et al, J Immunology 2000; 164: 3790].

Considering the lack of currently available treatment options for the majority of the conditions associated with protein kinases, especially GSK-3, Aurora-2, and Syk, there is still a great need for new therapeutic agents that inhibit these protein targets.

SUMMARY OF THE INVENTION

The present invention addresses this need by providing a compound of formula I:



I

or a pharmaceutically acceptable derivative thereof, wherein R¹ and Ar¹ are as defined below.

The present invention also provides a pharmaceutically acceptable composition comprising a compound of formula I.

The compounds and pharmaceutically acceptable compositions of the present invention are useful as

-7-

inhibitors of GSK-3, Aurora-2, and Syk protein kinases. Thus, they are also useful in methods for treating or lessening the severity of a variety of disorders, such as allergic diseases, proliferative disorders, cancer,
5 neurodegenerative disorders, and diabetes.

DESCRIPTION OF THE INVENTION

The present invention relates to a compound of formula I:



or a pharmaceutically acceptable derivative thereof, wherein:

R¹ is selected from R, halogen, CN, NO₂, or TR;

15 T is an optionally substituted C₁-C₄ alkylidene chain wherein up to two methylene units of T are optionally and independently replaced by O, N(R), C(O), S, SO, or SO₂;

each R is independently selected from hydrogen or an
20 optionally substituted C₁₋₆ aliphatic group, wherein:
two R bound to the same nitrogen atom are optionally taken together with the nitrogen to form a 3-7 membered saturated, partially unsaturated, or fully unsaturated ring having 0-2 heteroatoms, in
25 addition to the nitrogen bound thereto,
independently selected from nitrogen, oxygen, or sulfur;

Ar¹ is an optionally substituted ring selected from:

-8-

- (a) a 3-8 membered monocyclic or 8-10 membered bicyclic saturated, partially unsaturated, or aryl ring;
- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:
- Ar¹ is optionally substituted by one to four substituents selected from the group consisting of:
- (a) one group selected from QR, Ar², or QAr²;
- and
- (b) up to four R² groups;
- each Q is independently selected from a valence bond or an optionally substituted C₁₋₆ alkylidene chain, wherein: one or two non-adjacent methylene units of Q are optionally and independently replaced by -O-, -S-, -NR-, -C(O)-, -CO₂-, -C(O)NR-, -OC(O)NR-, -C(O)C(O)-, -C(O)C(O)-, -NRC(O)-, NRCO₂-, -NRC(O)NR-, -S(O)-, -SO₂-, -NRSO₂-, -SO₂NR-, or -NRSO₂NR-;
- each Ar² is an optionally substituted ring independently selected from:
- (a) a 3-8 membered monocyclic or 8-10 membered bicyclic saturated, partially unsaturated, or aryl ring;
- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or

-9-

(c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:

5 Ar² is optionally substituted by one to four R² groups; and

each R² is independently selected from R, halogen, NO₂, CN, OR, SR, N(R)₂, NRCOR, NRCON(R)₂, NRCO₂R, C(O)R, CO₂R, CON(R)₂, OC(O)N(R)₂, SOR, SO₂R, SO₂N(R)₂, NRSO₂R, NRSO₂N(R)₂, C(O)C(O)R, or C(O)CH₂C(O)R; wherein:

two R² on adjacent positions on Ar¹ or Ar² are optionally taken together to form a saturated, partially unsaturated, or fully unsaturated 4-6 membered ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

As used herein, the following definitions shall apply unless otherwise indicated.

10 The phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and each substitution is independent of the other.

15 The term "aliphatic" or "aliphatic group" as used herein means a straight-chain or branched C₁-C₈ hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic C₃-C₈ hydrocarbon or bicyclic C₈-C₁₂ hydrocarbon
20 that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic

-10-

ring system has 3-7 members. For example, suitable aliphatic groups include, but are not limited to, linear or branched or alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or
5 (cycloalkyl)alkenyl.

The terms "alkyl", "alkoxy", "hydroxyalkyl", "alkoxyalkyl", and "alkoxycarbonyl", used alone or as part of a larger moiety include both straight and branched chains containing one to twelve carbon atoms.
10 The terms "alkenyl" and "alkynyl" used alone or as part of a larger moiety shall include both straight and branched chains containing two to twelve carbon atoms.

The term "heteroatom" means nitrogen, oxygen, or sulfur and includes any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen.
15 Also the term "nitrogen" includes a substitutable nitrogen of a heterocyclic ring. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the
20 nitrogen may be N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl).

The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or
25 "aryloxyalkyl", refers to monocyclic, bicyclic and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term "aryl" may be used
30 interchangeably with the term "aryl ring". The term "aryl" also refers to heteroaryl ring systems as defined hereinbelow.

-11-

The term "heterocycle", "heterocyclyl", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic or tricyclic ring systems having five to fourteen ring members in which one or more ring members is a heteroatom, wherein each ring in the system contains 3 to 7 ring members.

The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents. Suitable substituents on the unsaturated carbon atom of an aryl, heteroaryl, aralkyl, or heteroaralkyl group are selected from halogen, R° , $-OR^\circ$, $-SR^\circ$, 1,2-methylene-dioxy, 1,2-ethylenedioxy, phenyl (Ph) optionally substituted with R° , $-O(Ph)$ optionally substituted with R° , $-CH_2(Ph)$ optionally substituted with R° , $-CH_2CH_2(Ph)$, optionally substituted with R° , $-NO_2$, $-CN$, $-N(R^\circ)_2$, $-NR^\circ C(O)R^\circ$, $-NR^\circ C(O)N(R^\circ)_2$, $-NR^\circ CO_2R^\circ$, $-NR^\circ NR^\circ C(O)R^\circ$, $-NR^\circ NR^\circ C(O)N(R^\circ)_2$, $-NR^\circ NR^\circ CO_2R^\circ$, $-C(O)C(O)R^\circ$, $-C(O)CH_2C(O)R^\circ$, $-CO_2R^\circ$, $-C(O)R^\circ$, $-C(O)N(R^\circ)_2$, $-OC(O)N(R^\circ)_2$, $-S(O)_2R^\circ$, $-SO_2N(R^\circ)_2$, $-S(O)R^\circ$, $-NR^\circ SO_2N(R^\circ)_2$, $-NR^\circ SO_2R^\circ$, $-C(=S)N(R^\circ)_2$, $-C(=NH)-N(R^\circ)_2$, or $-(CH_2)_yNHC(O)R^\circ$, wherein each R° is independently selected from hydrogen,

-12-

optionally substituted C₁₋₆ aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, -O(Ph), or -CH₂(Ph). Optional substituents on the aliphatic group of R^o are selected from NH₂, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo C₁₋₄ aliphatic.

An aliphatic group or a non-aromatic heterocyclic ring may contain one or more substituents. Suitable substituents on the saturated carbon of an aliphatic group or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and the following: =O, =S, =NNHR^{*}, =NN(R^{*})₂, =NNHC(O)R^{*}, =NNHCO₂(alkyl), =NNHSO₂(alkyl), or =NR^{*}, where each R^{*} is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic. Optional substituents on the aliphatic group of R^{*} are selected from NH₂, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic).

Optional substituents on the nitrogen of a non-aromatic heterocyclic ring are selected from -R^{*}, -N(R^{*})₂, -C(O)R^{*}, -CO₂R^{*}, -C(O)C(O)R^{*}, -C(O)CH₂C(O)R^{*}, -SO₂R^{*}, -SO₂N(R^{*})₂, -C(=S)N(R^{*})₂, -C(=NH)-N(R^{*})₂, or -NR^{*}SO₂R^{*}; wherein R^{*} is hydrogen, an optionally substituted C₁₋₆ aliphatic, optionally substituted phenyl, optionally substituted -O(Ph), optionally substituted -CH₂(Ph), optionally substituted -CH₂CH₂(Ph), or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring. Optional substituents on the aliphatic group or the phenyl ring of R^{*} are selected from NH₂, NH(C₁₋₄ aliphatic), N(C₁₋₄ aliphatic)₂, halogen, C₁₋₄ aliphatic, OH, O(C₁₋₄ aliphatic),

-13-

NO₂, CN, CO₂H, CO₂(C₁₋₄ aliphatic), O(halo C₁₋₄ aliphatic), or halo(C₁₋₄ aliphatic).

The term "alkylidene chain" refers to a straight or branched carbon chain that may be fully saturated or have one or more units of unsaturation and has two points of attachment to the rest of the molecule.


A combination of substituents or variables is permissible only if such a combination results in a stable or chemically feasible compound. A stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms, all such tautomeric forms of the compounds being within the scope of the invention.

Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

-14-


According to one embodiment, the present invention relates to a compound of formula I:



I

- 5 or a pharmaceutically acceptable derivative thereof,
wherein R¹ and Ar¹ are as defined above, provided that:
when Ar¹ is phenyl with two R² substituents, then the two
R² are not simultaneously OR in the meta and para
positions of Ar¹.

- 10 According to another embodiment, the present
invention relates to a compound of formula I:



I

- or a pharmaceutically acceptable derivative thereof,
15 wherein R¹ and Ar¹ are as defined above, provided that:
R¹ is other than CN.

Preferred Ar¹ groups of formula I are optionally substituted rings selected from:

- (a) a phenyl, indanyl, or naphthyl ring;
20 (b) a 5-6 membered heterocyclic ring having 1-3
heteroatoms independently selected from
nitrogen, oxygen, or sulfur; or
(c) a 5-6 membered monocyclic or 9-10 membered
bicyclic heteroaryl ring having 1-2 heteroatoms

-15-

independently selected from oxygen, nitrogen, or sulfur.

More preferred Ar¹ groups of formula I are rings selected from:

- 5 (a) a phenyl, indanyl, or naphthyl ring;
- (b) a 5-6 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic heteroaryl ring having
10 1-2 nitrogens, wherein:
- Ar¹ is substituted by one to four substituents selected from the group consisting of:
- (a) one group selected from QR, Ar², or QAr²; and
- 15 (b) up to four R² groups.

Most preferred Ar¹ rings are selected from substituted phenyl, indanyl, naphthyl, pyrimidinyl, or pyridyl. Preferred R² substituents on Ar¹ are halogen, CN, CO₂R, R, NO₂, OR, haloalkyl, SO₂N(R)₂, or N(R)₂. More
20 preferred R² substituents on Ar¹ are fluoro, iodo, chloro, bromo, CO₂CH₃, methyl, ethyl, t-butyl, NH₂, NHMe, N(Me)₂, OH, OCH₃, OCH₂CH₃, CF₃, SO₂NH₂, or SO₂NHMe. Other preferred compounds include those where two R² are taken together to form a methylenedioxy or an ethylenedioxy substituent.

25 Preferred QR or QAr² substituents on Ar¹ of formula I are those wherein Q is a C₁₋₄ alkylidene chain wherein one or two methylene units of Q are optionally replaced by O, NR, NRCO, NRCO₂, NRSO₂, or CONR, wherein each R is hydrogen or an optionally substituted C₁₋₄
30 aliphatic group and wherein Ar² is a 3-6 membered carbocyclic ring or an optionally substituted phenyl, 5-6 membered heterocyclic, or heteroaryl ring having one to

-16-

two heteroatoms independently selected from nitrogen, oxygen, or sulfur.

The Ar^2 group of the QAr^2 moiety of formula I is optionally substituted with R, OR, N(R)_2 , SO_2R , halogen, NO_2 , CN, SR, $\text{SO}_2\text{N(R)}_2$, CO_2R , C(O)R , or oxo. More preferred QAr^2 groups of formula I are selected from $\text{O(CH}_2)_3\text{pyrrolidin-1-yl}$, $\text{O(CH}_2)_2\text{morpholin-4-yl}$, $\text{O(CH}_2)_3(4\text{-hydroxyethylpiperazin-1-yl})$, $\text{O(CH}_2)_3\text{piperazin-1'-yl}$, $\text{O(CH}_2)_3(4\text{-hydroxymethylpiperidin-1-yl})$, $\text{O(CH}_2)_3(4\text{-hydroxypiperidin-1-yl})$, $\text{NHCOCH}_2\text{pyridin-2-yl}$, $\text{NHCOCH}_2(2\text{-aminothiazol-4-yl})$, $\text{NHCOCH}_2\text{cyclopropyl}$, $\text{NHCO(CH}_2)_2(\text{piperazin-2,5-dione-3-yl})$, $\text{NHCOpyrrolidin-1-yl}$, $\text{NHCOMorpholin-4-yl}$, $\text{NHCO}_2\text{CH}_2\text{tetrahydrofuran-2-yl}$, $\text{NHCO}_2\text{tetrahydrofuran-2-yl}$, $\text{NHCO}_2\text{tetrahydropyran-4-yl}$, $\text{NHCO}_2\text{CH}_2\text{tetrahydropyran-2-yl}$, Ophenyl, $\text{OCH}_2(\text{cyclohexyl})$, $\text{OCH}_2\text{phenyl}$, $\text{OCH}_2(3\text{-CN-phenyl})$, $\text{OCH}_2(2\text{-NO}_2\text{-phenyl})$, $\text{OCH}_2(3\text{-NH}_2\text{-phenyl})$, $\text{OCH}_2(4\text{-CO}_2\text{R-phenyl})$, $\text{OCH}_2\text{-pyridyl}$, $\text{OCH}_2(\text{mono-, di-, or tri-halogenated phenyl})$, $\text{OCH}_2(\text{C}_1\text{-C}_6\text{ aliphatic substituted phenyl})$, $\text{OCH}_2\text{CH}_2\text{-pyrrolyl}$, $\text{OCH}_2\text{-pyrrolyl}$, and $\text{OCH}_2(\text{phenyl substituted with one or two R}^2)$

Preferred Ar^2 substituents on Ar^1 of formula I are optionally substituted rings selected from:

- (a) a phenyl, indanyl, or naphthyl ring;
- (b) a 5-6 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 9-10 membered bicyclic heteroaryl ring having 1-2 heteroatoms independently selected from oxygen, nitrogen, or sulfur, wherein:


Ar^2 is optionally substituted with 1-2 R^2 groups.

-17-

More preferred Ar^2 substituents on Ar^1 of formula I are optionally substituted rings selected from phenyl, pyridyl, indolyl, naphthyl, or benzo[1,3]dioxolyl.

- 5 Preferred R^2 groups, when present, on the Ar^2 substituent on Ar^1 of formula I are selected from R, halogen, NO_2 , CN, OR, SR, $\text{N}(\text{R})_2$, $\text{C}(\text{O})\text{R}$, $\text{SO}_2\text{N}(\text{R})_2$, or SO_2R . More preferred R^2 groups, when present, on the Ar^2 substituent on Ar^1 of formula I are selected from methyl, 10 ethyl, t-butyl, fluoro, chloro, bromo, CF_3 , OMe, OEt, CN, SO_2Me , SO_2NH_2 , NH_2 , NHMe, $\text{N}(\text{Me})_2$, SMe, SEt, OH, $\text{C}(\text{O})\text{Me}$, NO_2 , or CH_2OH .

One embodiment of this invention relates to a compound of formula Ia:



15


Ia

or a pharmaceutically acceptable derivative thereof, wherein Ar^1 is as defined above.

- 20 Preferred Ar^1 groups of formula Ia are those described above for formula I.

Another embodiment of this invention relates to a compound of formula II:

-18-




II

or a pharmaceutically acceptable derivative thereof,
wherein R^2 is as defined above.

5 Preferred R^2 groups of formula II are those
defined above for formula I and include those where two R^2
are taken together to form a methylenedioxy or an
ethylenedioxy substituent.

A preferred embodiment of this invention
10 relates to a compound of formula IIa or IIa':




or a pharmaceutically acceptable derivative thereof,
wherein QR, QAr^2 and R^2 are as defined above.

15 Preferred R^2 groups of formula IIa or IIa' are
those described above for compounds of formula I.

Preferred QR and QAr^2 groups of formula IIa or
IIa' are those described above for compounds of formula
I.

20 Another preferred embodiment relates to a
compound of formula III:

-19-




III

or a pharmaceutically acceptable derivative thereof,
wherein R^1 , R^2 , and QAr^2 are as defined above.

5 Preferred R^2 and QAr^2 groups of formula III are
those described above for formula I.

Preferred R^1 groups of formula III are selected
from hydrogen, $N(R)_2$, SR, OR, or TR. More preferred R^1
groups are selected from OH, OCH_3 , CH_2OH , CH_2OCH_3 , CH_2NH_2 ,
10 or CH_2NHCH_3 .

Another preferred embodiment relates to a
compound of formula IV:



IV

15 or a pharmaceutically acceptable derivative thereof,
wherein R^2 and Ar^2 are as defined above.

Preferred R^2 and Ar^2 groups of formula IV are
those described above for compounds of formula I.

Exemplary structures of formula I are set forth
20 in Table 1 below.

Table 1. Compounds of Formula I

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
1		2		3	
4		5		6	
7		8		9	
10		11		12	
13		14		15	
16		17		18	

- 21 -

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
19		20		21	
22		23		24	
25		26		27	
28		29		30	
31		-	-	33	
34		35		36	
37		38		39	

- 22 -

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
40		41		42	
43		44		45	
46		47		48	
49		50		51	
52		53		54	
55		56		57	
58		59		60	

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
61		62		63	
64		65		66	
67		68		69	
70		71		72	
73		74		75	
76		77		78	
79		80		81	



Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
82		83		84	
85		86		87	
88		89		90	
91		92		93	
94		95		96	
97		98		99	
100		101		-	-

- 25 -


Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
103		104		105	
106		107		108	
109		110		111	
112		113		114	
115		116		117	

Exemplary structures of formulae IIa and IIa' are set forth in Table 2 below.


5 Table 2. Compounds of Formulae IIa and IIa'




- 26 -




IIa-3




IIa-4




IIa-5




IIa-6




IIa-7




IIa-8




IIa-9




IIa-10




IIa-11




IIa-12




IIa-13



IIa-14




IIa-15




IIa-16


-27-




IIa-17




IIa-18




IIa-19




IIa-20




IIa-21




IIa-22




IIa-23




IIa-24




IIa-25



IIa-26




IIa-27




IIa-28


-28-




IIa-29




IIa-30




IIa-31




IIa-32




IIa-33




IIa-34




IIa-35




IIa-36




IIa-37



IIa-38




IIa-39




IIa-40


-29-




IIa-41




IIa-42




IIa-43




IIa-44




IIa-45




IIa-46




IIa-47



IIa-48



IIa-49




IIa-50

The present compounds may be prepared in general by methods known to those skilled in the art for analogous compounds, as illustrated by the general Scheme I and the synthetic examples shown below.

-30-

Scheme I



Reagents and conditions: (a) DMF-DMA, THF, 12-18 hours, room temperature; (b) H_2NCN , 4N HCl in dioxane, 12-18 hours, 120°C; (c) Ethanol, reflux, 12-18 hours.

Scheme I above shows a general synthetic route that may be used for preparing compounds of formula I.

In step (a), a solution of 2-acetyl thiazole (1) in THF is treated with dimethylformamide-dimethylacetal and the resulting mixture stirred at room temperature over night. The reaction mixture is concentrated *in vacuo* and the concentrate triturated with diethyl ether to afford 2.

To prepare intermediate 4 from aniline 3 in step (b), a mixture of 3 and cyanamide in HCl (4N in dioxane) is heated at 120°C overnight. After cooling to room temperature, aqueous work-up affords the desired guanidine compound 4. One of skill in the art would recognize that a wide variety of aryl guanidines may be prepared at step (b) and, thus, be used to prepare compounds of formula I with a wide variety of Ar^1 rings.

In step (c), guanidine 4 is combined with enaminone 2 in ethanol in a sealed tube. The resulting mixture is heated at reflux overnight then concentrated and the crude product purified by column chromatography

-31-

to afford the desired pyrimidine compound 5. The details of the conditions used for producing these compounds are set forth in the Examples.

The activity of a compound utilized in this invention as an inhibitor of GSK3, Aurora2, or Syk protein kinase may be assayed *in vitro*, *in vivo* or in a cell line according to methods known in the art and by the methods set forth in the Examples below. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity or ATPase activity of activated GSK3, Aurora2, or Syk protein kinase. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to GSK3, Aurora2, or Syk protein kinase. Inhibitor binding may be measured by radiolabelling the inhibitor prior to binding, isolating the inhibitor/GSK3, inhibitor/Aurora2, or inhibitor/Syk complex and determining the amount of radiolabel bound. Alternatively, inhibitor binding may be determined by running a competition experiment where new inhibitors are incubated with GSK3, Aurora2, or Syk protein kinase bound to known radioligands. Detailed conditions for assaying a compound utilized in this invention as an inhibitor of GSK3, Aurora2, or Syk protein kinase are set forth in the Examples below.

According to another embodiment, the present invention provides a composition, or pharmaceutically acceptable composition, comprising a compound of this invention or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier, adjuvant, or vehicle. The amount of compound in the compositions of this invention is such that is effective to detectably inhibit a protein kinase, particularly GSK3, Aurora2, or Syk protein kinase, in a biological

-32-

sample or in a patient. Preferably the composition of this invention is formulated for administration to a patient in need of such composition. Most preferably, the composition of this invention is formulated for oral
5 administration to a patient.

The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the
10 pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this invention include, but are not
15 limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or
20 electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium
25 carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

The term "detectably inhibit", as used herein means a measurable change in GSK3, Aurora2, or Syk
30 protein kinase activity between a sample comprising said composition and a GSK3, Aurora2, or Syk protein kinase and an equivalent sample comprising GSK3, Aurora2, or Syk protein kinase in the absence of said composition.

-33-

A "pharmaceutically acceptable derivative" means any non-toxic salt, ester, salt of an ester or other derivative of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of a GSK3, Aurora2, or Syk protein kinase.

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth

-34-

metal (e.g., magnesium), ammonium and $N^+(C_{1-4} \text{ alkyl})_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible
5 products may be obtained by such quaternization.

The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as
10 used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally,
15 intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and
20 suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be
25 employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

For this purpose, any bland fixed oil may be
30 employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as

-35-

olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

Alternatively, the pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

-36-

The pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic

-37-

uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

The pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

Most preferably, the pharmaceutically acceptable compositions of this invention are formulated for oral administration.

The amount of the compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

-38-

Depending upon the particular condition, or disease, to be treated or prevented, additional therapeutic agents, which are normally administered to treat or prevent that condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

For example, in the treatment of diabetes other anti-diabetic agents may be combined with the compounds of this invention to treat diabetes. These agents include, without limitation, insulin in injectable or inhalation form, insulin analogues, glitazones, sulfonyl ureas, alpha glucosidase inhibitors, biguanides, and insulin sensitizers.

Other examples of agents the inhibitors of this invention may also be combined with include, without limitation: chemotherapeutic agents such as Gleevec™, adriamycin, dexamethasone, vincristine, cyclophosphamide, fluorouracil, topotecan, taxol, interferons, and platinum derivatives; treatments for Alzheimer's Disease such as Aricept® and Exelon®; treatments for Parkinson's Disease such as L-DOPA/carbidopa, entacapone, ropinrole, pramipexole, bromocriptine, pergolide, trihexephendyl, and amantadine; agents for treating Multiple Sclerosis (MS) such as beta interferon (e.g., Avonex® and Rebif®), Copaxone®, and mitoxantrone; treatments for asthma such as albuterol and Singulair®; agents for treating schizophrenia such as zyprexa, risperdal, seroquel, and haloperidol; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons,

-39-

corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-
5 Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for
10 treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; and agents for treating immunodeficiency disorders such as gamma globulin.

The amount of additional therapeutic agent present in the compositions of this invention will be no
15 more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally
20 present in a composition comprising that agent as the only therapeutically active agent.

According to another embodiment, the invention relates to a method of inhibiting GSK3 protein kinase activity in a biological sample comprising the step of
25 contacting said biological sample with a compound of this invention, or a composition comprising said compound.

According to another embodiment, the invention relates to a method of inhibiting Aurora2 protein kinase activity in a biological sample comprising the step of
30 contacting said biological sample with a compound of this invention, or a composition comprising said compound.

According to another embodiment, the invention relates to a method of inhibiting Syk protein kinase

-40-

activity in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a composition comprising said compound.

The term "biological sample", as used herein,
5 includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

Inhibition of GSK3, Aurora2, or Syk protein
10 kinase activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, biological specimen storage, and biological assays.

15 According to another embodiment, the invention provides a method for treating or lessening the severity of a GSK3-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to the present invention.

20 The term "GSK3-mediated disease" or "GSK3-mediated condition", as used herein, means any disease or other deleterious condition in which GSK3 protein kinase is known to play a role. Such conditions include, without limitation, diabetes, neurodegenerative
25 disorders, Alzheimer's disease, Huntington's, Parkinson's, AIDS associated dementia, amyotrophic lateral sclerosis (ALS), multiple sclerosis (MS), schizophrenia, stroke, cardiomyocyte hypertrophy, and baldness.

30 According to another embodiment, the invention provides a method for treating or lessening the severity of an Aurora2-mediated disease or condition in a patient

-41-

comprising the step of administering to said patient a composition according to the present invention.

The term "Aurora2-mediated disease" or "Aurora2-mediated condition", as used herein, means any
5 disease or other deleterious condition in which Aurora2 protein kinase is known to play a role. Such conditions include, without limitation, cancers such as colon and breast cancer.

According to another embodiment, the invention
10 provides a method for treating or lessening the severity of a Syk-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to the present invention.


The term "Syk-mediated disease" or "Syk-mediated condition", as used herein, means any disease or
15 other deleterious condition in which Syk protein kinase is known to play a role. Such conditions include, without limitation, allergic disorders, especially asthma.

20 In an alternate embodiment, the methods of this invention that utilize compositions that do not contain an additional therapeutic agent, comprise the additional step of separately administering to said patient an additional therapeutic agent. When these additional
25 therapeutic agents are administered separately they may be administered to the patient prior to, sequentially with or following administration of the compositions of this invention.

In order that the invention described herein
30 may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLES

Example 1




2

3-Dimethylamino-1-thiazol-2-yl-propenone (2): To a solution of 2-acetylthiazole (500 mg, 3.93 mmol) in 2 mL of THF was added dimethylformamide-dimethylacetal (1.6 ml, 7.86 mmol) and the resulting solution was stirred at room temperature overnight. The reaction mixture was evaporated and the residue triturated with ethyl acetate and the product isolated by filtration. The filtered solid was washed with diethyl ether and dried to afford 2 (400 mg) as a yellow solid.

15

Example 2




4

N-(3-Benzyloxy-phenyl)-guanidine (4): 3-benzyloxy aniline
20 (1 g, 5 mmol) and cyanamide (420 mg, 10 mmol) were
dissolved in 4 mL of 4N HCl in dioxane and heated to 120°C
overnight. The reaction mixture was cooled to room
temperature, quenched with water, and extracted with
diethyl ether. The aqueous layer was made basic with 2N
25 NaOH and extracted thrice with dichloromethane. The
organic extracts were combined and washed with brine,

-43-

dried over MgSO_4 , and concentrated to afford 4 (450 mg) as a light brown solid.

Example 3



I-7

(3-Benzyloxy-phenyl)-(4-thiazol-2-yl-pyrimidin-2-yl)-amine (I-7): A solution of 3-benzyloxy guanidine 4 (50 mg, 0.20 mmol) and enaminone 2 (50 mg, 0.27 mmol) in ethanol (3 mL) was heated at reflux in a sealed tube overnight. The reaction mixture was concentrated and the residue purified by column chromatography to afford compound I-7 (20 mg). $M+1$ (obs) 347.1. $R_t = 4.28$ minutes.

Example 4

We have prepared other compounds of formula I by methods substantially similar to those described in the above Examples 1-3 and those illustrated in Scheme I. The characterization data for these compounds is summarized in Table 3 below and includes LC/MS ($M+1$ observed), HPLC, and ^1H NMR data.

The term " $R_t(\text{min})$ " refers to the retention time, in minutes, associated with the compound using the following HPLC method. The R_t was determined using a Waters ODS-AQ (2 x 50mm) column at ambient temperature with a gradient of 10-90% CH_3CN in water over 5 minutes and with the absorbance detected as an average of 190-380 nm @ 4 nm increments.

-44-

The term "Y" designates that ^1H NMR data was obtained and found to be consistent with the assigned structure. Compound numbers correspond to the compound numbers listed in Table 1.

5

Table 3. Characterization Data for Selected Compounds of Formula I

Compound No	M+1 (obs)	R _t (min)	^1H NMR
I-16	300.0	3.15	-
I-19	273.0	3.18	-
I-22	289.0	3.47	-
I-28	290.0	2.76	-
I-31	381.0	3.66	-
I-34	334.8	3.54	-
I-37	323.0	3.56	-
I-40	300.0	3.19	-
I-43	273.0	3.1	-
I-46	289.0	3.19	-
I-49	291.0	3.33	-
I-52	305.0	3.62	-
I-55	334.0	2.16	-
I-58	309.0	3.60	-
I-61	322.9	2.99	-
I-64	291.0	2.67	-
I-67	269.0	3.28	-
I-73	283.0	3.53	-
I-10	361.1	3.76	-
I-13	347.0	3.82	-
I-4	347.1	4.14	-
I-7	361.1	4.28	-
I-25	285.0	2.98	-
I-76	367.2	4.53	-
I-79	386.1	3.58	-
I-82	419.1	3.80	-
I-85	397.1	3.94	-
I-88	395.1	4.11	-
I-100	379.1	3.85	-

- 45 -

Compound No	M+1 (obs)	R _t (min)	¹ H NMR
I-103	397.1	3.98	-
I-106	429.1	4.13	-
I-109	379.1	3.84	-
I-112	429.0	4.40	-
I-115	375.2	3.96	-
I-2	429.0	4.38	-
I-5	439.1	3.24	-
I-8	362.1	2.21	-
I-11	-	-	Y
I-14	376.2	2.56	-
I-17	405.1	3.72	-
I-20	362.1	2.54	-
I-23	362.1	2.55	-
I-29	364.2	3.52	-
I-35	406.1	3.8	Y
I-38	376.1	3.13	Y
I-41	379.2	3.84	-
I-44	391.1	3.84	-
I-47	429.1	4.12	-
I-91	429.1	4.12	-
I-94	397.1	3.75	-

Example 5GSK-3 Inhibition Assay

Compounds were screened for their ability to

5 inhibit GSK3- β (AA 1-420) activity using a standard coupled enzyme system (Fox et al. (1998) *Protein Sci.* 7, 2249). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 300 μ M NADH, 1 mM DTT and 1.5% DMSO. Final substrate

10 concentrations in the assay were 10 μ M ATP (Sigma Chemicals, St Louis, MO) and 300 μ M peptide (HSSPHQS(PO₃H₂)EDEEE, American Peptide, Sunnyvale, CA). Reactions were carried out at 30°C and 60 nM GSK-3 β .

-46-

Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 300 μ M NADH, 30 μ g/ml pyruvate kinase and 10 μ g/ml lactate dehydrogenase.

5 An assay stock buffer solution was prepared containing all of the reagents listed above with the exception of ATP and the test compound of interest. 59 μ l of the test reaction was placed in a 96 well $\frac{1}{2}$ -diameter plate (Corning, Corning, NY) then treated with 1
10 μ l of a 2 mM DMSO stock containing the test compound (final compound concentration 30 μ M). The plate was incubated for ~10 minutes at 30 °C then the reaction initiated by addition of 7 μ l of ATP (final concentration 10 μ M). Rates of reaction were obtained using a
15 Molecular Devices Spectramax plate reader (Sunnyvale, CA) over a 5 minute read time at 30 °C. Compounds showing greater than 50 % inhibition versus standard wells containing DMSO, but no compound, were titrated and K_i values were determined.

20 Table 4 shows the results of the activity of selected compounds of this invention in the GSK3 inhibition assay. The compound numbers correspond to the compound numbers in Table 1. Compounds having a K_i less than 0.1 micromolar (μ M) are rated "A", compounds having
25 a K_i between 0.1 and 1 μ M are rated "B" and compounds having a K_i greater than 1 μ M are rated "C".

Table 4. GSK3- β Activity of Selected Compounds

No. I-	Activity	No. I-	Activity	No. I-	Activity
16	C	19	A	22	A
28	A	31	A	34	A
37	A	40	A	43	A

-47-

No. I-	Activity	No. I-	Activity	No. I-	Activity
46	A	49	A	52	A
55	A	58	A	61	C
64	C	67	B	73	A
10	C	4	A	7	-
25	A	76	A	79	A
82	A	85	A	88	A
100	A	103	A	106	A
109	A	112	A	115	A
2	C	5	A	8	A
11	A	14	A	17	A
20	A	23	A	29	A
35	A	38	A	41	A
44	A	47	A	91	A
94	A	3	A	6	A
9	A	12	A	15	A
18	A	21	C	24	A
27	A	30	A	33	A
36	B	39	A	42	A
45	A	48	A	51	B
54	A	57	A	60	A
63	A	66	A	69	A
72	A	75	A	78	A
81	A	84	A	87	A
90	A	93	C	96	C
99	A	-	-	105	A
108	B	111	A	114	C
117	C	-	-	-	-

Example 6Aurora2 Inhibition Assay

Compounds are screened in the following manner

5 for their ability to inhibit Aurora2 using a standard coupled enzyme assay (Fox et al (1998) *Protein Sci* 7, 2249). To an assay stock buffer solution containing 0.1M HEPES 7.5, 10 mM MgCl₂, 1 mM DTT, 25 mM NaCl, 2.5 mM

-48-

phosphoenolpyruvate, 300 mM NADH, 30 mg/ml pyruvate kinase, 10 mg/ml lactate dehydrogenase, 40 mM ATP, and 800 μ M peptide (LRRASLG, American Peptide, Sunnyvale, CA) is added a DMSO solution of a compound of the present invention to a final concentration of 30 μ M. The resulting mixture is incubated at 30 °C for 10 minutes. The reaction was initiated by the addition of 10 μ L of Aurora2 stock solution to give a final concentration of 70 nM in the assay. The rates of reaction are obtained by monitoring absorbance at 340 nm over a 5 minute read time at 30 °C using a BioRad Ultramark plate reader (Hercules, CA). The IC₅₀ values are determined from the rate data as a function of inhibitor concentration.

Table 5 shows the results of the activity of selected compounds of this invention in the Aurora2 inhibition assay. The compound numbers correspond to the compound numbers in Table 1. Compounds having an IC₅₀ less than 0.5 micromolar (μ M) are rated "A", compounds having an IC₅₀ between 0.5 and 2 μ M are rated "B" and compounds having an IC₅₀ greater than 2 μ M are rated "C".

Table 5. Aurora2 Activity of Selected Compounds

No. I-	Activity	No. I-	Activity	No. I-	Activity
2	C	4	B	5	C
7	C	8	A	10	C
11	A	13	C	14	B
16	A	17	A	19	B
20	A	22	A	23	A
25	B	26	A	28	C
29	A	31	A	-	-
34	A	35	C	37	B
38	A	40	C	41	C
43	C	46	C	49	B
50	B	52	C	55	A

-49-

No. I-	Activity	No. I-	Activity	No. I-	Activity
56	B	58	C	59	C
61	C	62	B	64	C
67	A	70	B	73	A
76	C	79	C	82	C
85	C	88	B	91	C
94	A	97	C	100	C
103	C	106	C	109	C
112	C	114	B	-	-

Example 7

Syk Inhibition Assay

Compounds were screened for their ability to
5 inhibit Syk using a standard coupled enzyme assay (Fox et
al (1998) *Protein Sci* 7, 2249). Reactions were carried
out in 100 mM HEPES pH 7.5, 10 mM MgCl₂, 25 mM NaCl, 1 mM
DTT and 1.5% DMSO. Final substrate concentrations in the
assay were 200 μ M ATP (Sigma chemical Co.) and 4 μ M poly
10 Gly-Tyr peptide (Sigma Chemical Co.). Assays were
carried out at 30 °C and 200 nM Syk. Final
concentrations of the components of the coupled enzyme
system were 2.5 mM phosphoenolpyruvate, 300 μ M NADH, 30
 μ g/ml pyruvate kinase and 10 μ g/ml lactate dehydrogenase.

15 An assay stock buffer solution was prepared
containing all of the reagents listed above, with the
exception of Syk, DTT and the test compound of interest.
56 μ l of the test reaction was placed in a 96 well plate
followed by the addition of 1 μ l of 2 mM DMSO stock
20 containing the test compound (final compound
concentration 30 μ M). The plate was pre-incubated for
~10 minutes at 30 °C and the reaction initiated by the
addition of 10 μ l of enzyme (final concentration 25 nM).
Rates of reaction were obtained using a BioRad Ultramark
25 plate reader (Hercules, CA) over a 5 minute read time at

-50-

30°C. Compounds showing >50 % inhibition versus standard wells containing DMSO, but no compound, were titrated and IC₅₀'s determined using a similar protocol.

Table 6 shows the results of the activity of selected compounds of this invention in the Syk inhibition assay. The compound numbers correspond to the compound numbers in Table 1. Compounds having an IC₅₀ less than 0.5 micromolar (μ M) are rated "A", compounds having an IC₅₀ between 0.5 and 2 μ M are rated "B" and compounds having an IC₅₀ greater than 2 μ M are rated "C".

Table 6. Syk Activity of Selected Compounds

No. I-	Activity	No. I-	Activity	No. I-	Activity
2	C	4	B	5	A
7	A	8	C	11	A
14	B	16	B	17	A
19	B	22	A	23	A
25	A	29	A	31	A
34	A	35	C	37	A
38	A	43	C	44	A
46	C	47	C	49	C
50	A	52	C	53	B
55	A	56	B	58	C
59	C	61	C	62	A
64	C	67	A	73	A
76	C	82	C	88	C
91	C	94	A	106	C
109	C	112	C	115	C

While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention


-51-

is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example.

-52-

CLAIMSWe claim:

1. A compound of formula I:



I

or a pharmaceutically acceptable derivative thereof,
wherein:

R¹ is selected from R, halogen, CN, NO₂, or TR;

T is an optionally substituted C₁-C₄ alkylidene chain
wherein up to two methylene units of T are optionally
and independently replaced by O, N(R), C(O), S, SO, or
SO₂;

each R is independently selected from hydrogen or an
optionally substituted C₁₋₆ aliphatic group, wherein:

two R bound to the same nitrogen atom are optionally
taken together with the nitrogen to form a 3-7
membered saturated, partially unsaturated, or
fully unsaturated ring having 0-2 heteroatoms, in
addition to the nitrogen bound thereto,
independently selected from nitrogen, oxygen, or
sulfur;

Ar¹ is an optionally substituted ring selected from:

(a) a 3-8 membered monocyclic or 8-10 membered
bicyclic saturated, partially unsaturated, or
aryl ring;

-53-

- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:

Ar¹ is optionally substituted by one to four substituents selected from the group consisting of:

(a) one group selected from QR, Ar², or QAr²;

5

and

(b) up to four R² groups;

each Q is independently selected from a valence bond or an optionally substituted C₁₋₆ alkylidene chain, wherein:

one or two non-adjacent methylene units of Q are optionally and independently replaced by -O-, -S-, -NR-, -C(O)-, -CO₂-, -C(O)NR-, -OC(O)NR-, -C(O)C(O)-, -NRC(O)-, NRCO₂-, -NRC(O)NR-, -S(O)-, -SO₂-, -NRSO₂-, -SO₂NR-, or -NRSO₂NR-;

each Ar² is an optionally substituted ring independently selected from::

- (a) a 3-8 membered monocyclic or 8-10 membered bicyclic saturated, partially unsaturated, or aryl ring;
- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:

-54-

Ar² is optionally substituted by one to four R² groups; and

each R² is independently selected from R, halogen, NO₂, CN, OR, SR, N(R)₂, NRCOR, NRCON(R)₂, NRCO₂R, C(O)R, CO₂R, CON(R)₂, OC(O)N(R)₂, SOR, SO₂R, SO₂N(R)₂, NRSO₂R, NRSO₂N(R)₂, C(O)C(O)R, or C(O)CH₂C(O)R; wherein:

two R² on adjacent positions on Ar¹ or Ar² are optionally taken together to form a saturated, partially unsaturated, or fully unsaturated 4-6 membered ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

provided that:

- (a) when Ar¹ is phenyl, two R² are not simultaneously OR in the meta and para positions of Ar¹; and
- (b) R¹ is other than CN.

2. The compound according to claim 1, wherein Ar¹ is a ring selected from:

- (a) a phenyl, indanyl, or naphthyl ring;
- (b) a 5-6 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic heteroaryl ring having 1-2 nitrogens, wherein:

Ar¹ is substituted by one to four substituents selected from the group consisting of:

- (a) one group selected from QR, Ar², or QAr²;
- and
- (b) up to four R² groups.

3. The compound according to claim 2, wherein: Ar¹ is selected from a phenyl, indanyl, naphthyl, pyrimidinyl, or pyridyl ring.

-55-

4. The compound according to either of claims 2 or 3, wherein:

Ar¹ is substituted by one to four substituents selected from the group consisting of:

(a) one group selected from QR or QAr²; and

(b) up to four R² groups; wherein:

each R² is independently selected from halogen, CN, CO₂R, R, NO₂, OR, haloalkyl, SO₂N(R)₂, or N(R)₂;

each Q is independently selected from a C₁₋₄ alkylidene chain wherein one or two methylene units of Q are optionally replaced by O, NH, NHCO, NHCO₂, NHSO₂, or CONH; and

Ar² is a 3-6 membered carbocyclic ring or an optionally substituted phenyl or 5-6 membered heterocyclic or heteroaryl ring having one to two heteroatoms independently selected from nitrogen, oxygen, or sulfur.

5. The compound according to either of claims 2 or 3, wherein:

Ar¹ is substituted by Ar² and optionally substituted with 1-2 R² substituents, wherein:

each Ar² is an optionally substituted ring independently selected from:

(a) a phenyl, indanyl, or naphthyl ring;

(b) a 5-6 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or

(c) a 5-6 membered monocyclic or 9-10 membered bicyclic heteroaryl ring having 1-2 heteroatoms

-56-

independently selected from oxygen, nitrogen,
or sulfur, wherein:

Ar² is optionally substituted with 1-2 R²
groups; and

5 each R² is independently selected from R, halogen,
NO₂, CN, OR, SR, N(R)₂, C(O)R, SO₂N(R)₂, or SO₂R.

6. A compound selected from the following Table 1
compounds:

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
-	-	2		3	
4		5		6	
7		8		9	
10		11		12	
13		14		15	

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
16		17		18	
19		20		21	
22		23		24	
25		26		27	
28		29		30	
31				33	
34		35		36	

- 58 -

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
37		38		39	
40		41		42	
43		44		45	
46		47		48	
49		50		51	
52		53		54	
55		56		57	

Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
58		59		60	
61		62		63	
64		65		66	
67		68		69	
70		71		72	
73		74		75	
76		77		78	

- 60 -


Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
79		80		81	
82		83		84	
85		86		87	
88		89		90	
91		92		93	
94		95		96	
97		98		99	

-61-


Cmpd I-	Structure	Cmpd I-	Structure	Cmpd I-	Structure
100		101		-	-
103		104		105	
106		107		108	
109		110		111	
112		113		114	
115		116		117	

7. A compound selected from the following Table 2 compounds:


- 62 -




IIa-1




IIa-2




IIa-3




IIa-4




IIa-5




IIa-6




IIa-7




IIa-8




IIa-9




IIa-10




IIa-11



IIa-12




IIa-13




IIa-14


- 63 -




IIa-15




IIa-16




IIa-17




IIa-18




IIa-19




IIa-20




IIa-21




IIa-22




IIa-23



IIa-24




IIa-25




IIa-26


- 64 -




IIa-27




IIa-28




IIa-29




IIa-30




IIa-31




IIa-32




IIa-33




IIa-34




IIa-35



IIa-36




IIa-37




IIa-38


- 65 -




IIa-39




IIa-40




IIa-41




IIa-42




IIa-43




IIa-44




IIa-45




IIa-46




IIa-47



IIa-48



IIa-49



IIa-50


-66-

8. A composition comprising a compound according to claim 1, in an amount to detectably inhibit GSK3, Aurora2, or Syk protein kinase activity, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

9. The composition according to claim 8, additionally comprising an additional therapeutic agent selected from an anti-diabetic agent, a chemotherapeutic or anti-proliferative agent, a treatment for Alzheimer's Disease, a treatment for Parkinson's Disease, an agent for treating Multiple Sclerosis (MS), a treatment for asthma, an agent for treating schizophrenia, an anti-inflammatory agent, an immunomodulatory or immunosuppressive agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating liver disease, an agent for treating a blood disorder, or an agent for treating an immunodeficiency disorder.

10. A method of inhibiting GSK3, Aurora2, or Syk kinase activity in a biological sample, comprising the step of contacting said biological sample with:

- a) a composition according to claim 8; or
- b) a compound of formula I:



I

or a pharmaceutically acceptable derivative thereof, wherein:

R¹ is selected from R, halogen, CN, NO₂, or TR;

-67-

T is an optionally substituted C₁-C₄ alkylidene chain wherein up to two methylene units of T are optionally and independently replaced by O, N(R), C(O), S, SO, or SO₂;

each R is independently selected from hydrogen or an optionally substituted C₁₋₆ aliphatic group, wherein: two R bound to the same nitrogen atom are optionally taken together with the nitrogen to form a 3-7 membered saturated, partially unsaturated, or fully unsaturated ring having 0-2 heteroatoms, in addition to the nitrogen bound thereto, independently selected from nitrogen, oxygen, or sulfur;

Ar¹ is an optionally substituted ring selected from:

- (a) a 3-8 membered monocyclic or 8-10 membered bicyclic saturated, partially unsaturated, or aryl ring;
- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:

Ar¹ is optionally substituted by one to four substituents selected from:

- (a) one group selected from QR, Ar², or QAr²;
- and

- 5 (b) up to four R² groups;

each Q is independently selected from a valence bond or an optionally substituted C₁₋₆ alkylidene chain, wherein:

-68-

one or two non-adjacent methylene units of Q are optionally and independently replaced by -O-, -S-, -NR-, -C(O)-, -CO₂-, -C(O)NR-, -OC(O)NR-, -C(O)C(O)-, -NRC(O)-, NRCO₂-, -NRC(O)NR-, -S(O)-, -SO₂-, -NRSO₂-, -SO₂NR-, or -NRSO₂NR-;

each Ar² is an optionally substituted ring independently selected from::

- (a) a 3-8 membered monocyclic or 8-10 membered bicyclic saturated, partially unsaturated, or aryl ring;
- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:

Ar² is optionally substituted by one to four R² groups; and


each R² is independently selected from R, halogen, NO₂, CN, OR, SR, N(R)₂, NRCOR, NRCON(R)₂, NRCO₂R, C(O)R, CO₂R, CON(R)₂, OC(O)N(R)₂, SOR, SO₂R, SO₂N(R)₂, NRSO₂R, NRSO₂N(R)₂, C(O)C(O)R, or C(O)CH₂C(O)R; wherein:

two R² on adjacent positions on Ar¹ or Ar² are optionally taken together to form a saturated, partially unsaturated, or fully unsaturated 4-6 membered ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

11. A method of treating or lessening the severity of a GSK3-, Aurora2-, or Syk-mediated disease or condition in a patient, comprising the step of administering to said patient:

-69-

- a) a composition according to claim 8; or
- b) a compound of formula I:



I

or a pharmaceutically acceptable derivative thereof,
wherein:

- R¹ is selected from R, halogen, CN, NO₂, or TR;
- T is an optionally substituted C₁-C₄ alkylidene chain
wherein up to two methylene units of T are optionally
and independently replaced by O, N(R), C(O), S, SO, or
SO₂;
- each R is independently selected from hydrogen or an
optionally substituted C₁₋₆ aliphatic group, wherein:
two R bound to the same nitrogen atom are optionally
taken together with the nitrogen to form a 3-7
membered saturated, partially unsaturated, or
fully unsaturated ring having 0-2 heteroatoms, in
addition to the nitrogen bound thereto,
independently selected from nitrogen, oxygen, or
sulfur;
- Ar¹ is an optionally substituted ring selected from:
 - (a) a 3-8 membered monocyclic or 8-10 membered
bicyclic saturated, partially unsaturated, or
aryl ring;
 - (b) a 3-7 membered heterocyclic ring having 1-3
heteroatoms independently selected from
nitrogen, oxygen, or sulfur; or
 - (c) a 5-6 membered monocyclic or 8-10 membered
bicyclic heteroaryl ring having 1-4 heteroatoms

-70-

independently selected from nitrogen, oxygen, or sulfur, wherein:

Ar¹ is optionally substituted by one to four substituents selected from:

(a) one group selected from QR, Ar², or QAr²; and

5 (b) up to four R² groups;

each Q is independently selected from a valence bond or an optionally substituted C₁₋₆ alkylidene chain, wherein:

one or two non-adjacent methylene units of Q are optionally and independently replaced by -O-, -S-, -NR-, -C(O)-, -CO₂-, -C(O)NR-, -OC(O)NR-, -C(O)C(O)-, -NRC(O)-, NRCO₂-, -NRC(O)NR-, -S(O)-, -SO₂-, -NRSO₂-, -SO₂NR-, or -NRSO₂NR-;

each Ar² is an optionally substituted ring independently selected from::

- (a) a 3-8 membered monocyclic or 8-10 membered bicyclic saturated, partially unsaturated, or aryl ring;
- (b) a 3-7 membered heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5-6 membered monocyclic or 8-10 membered bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein:

Ar² is optionally substituted by one to four R² groups; and

each R² is independently selected from R, halogen, NO₂, CN, OR, SR, N(R)₂, NRCOR, NRCON(R)₂, NRCO₂R, C(O)R, CO₂R, CON(R)₂, OC(O)N(R)₂, SOR, SO₂R, SO₂N(R)₂, NRSO₂R, NRSO₂N(R)₂, C(O)C(O)R, or C(O)CH₂C(O)R; wherein:

-71-

two R² on adjacent positions on Ar¹ or Ar² are optionally taken together to form a saturated, partially unsaturated, or fully unsaturated 4-6 membered ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

12. The method according to claim 11, wherein said GSK3-mediated disease is selected from diabetes, a neurodegenerative disorder, a CNS disorder, Alzheimer's disease, Huntington's, Parkinson's, AIDS associated dementia, amyotrophic lateral sclerosis (AML), multiple sclerosis (MS), stroke, schizophrenia, cardiomyocyte hypertrophy, a manic depressive disorder, or baldness.

13. The method according to claim 11, wherein said Aurora2-mediated disease is selected from cancer.

14. The method according to claim 11, wherein said Syk-mediated disease is selected from an allergic disorder.

15. The method according to claim 11, comprising the additional step of administering to said patient an additional therapeutic agent selected from an anti-diabetic agent, a chemotherapeutic or anti-proliferative agent, a treatment for Alzheimer's Disease, a treatment for Parkinson's Disease, an agent for treating Multiple Sclerosis (MS), a treatment for asthma, an agent for treating schizophrenia, an anti-inflammatory agent, an immunomodulatory or immunosuppressive agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating liver disease, an agent

-72-

for treating a blood disorder, or an agent for treating an immunodeficiency disorder, wherein:

said additional therapeutic agent is appropriate for the disease being treated; and

said additional therapeutic agent is administered together with said composition as a single dosage form or separately from said composition as part of a multiple dosage form.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 02/16352

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/04 A61K31/427 A61P03/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 958 935 A (DAVIS JEREMY MARTIN ET AL) 28 September 1999 (1999-09-28) claims 1,20,21; example 8	1-15
X	WO 00 78731 A (CELLTECH CHIROSCIENCE LTD ;DAVIS JEREMY MARTIN (GB); MOFFAT DAVID) 28 December 2000 (2000-12-28) page 2, line 12 - line 14; claim 1; examples 50-52,54,55,57	1-15



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

1 August 2002

Date of mailing of the international search report

09/08/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seelmann, I

INTERNATIONAL SEARCH REPORT
 Information on patent family members

International Application No
 PCT/US 02/16352

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
US 5958935	A	28-09-1999	AU	7631496 A		11-06-1997
			EP	0862560 A1		09-09-1998
			WO	9719065 A1		29-05-1997
			US	6235746 B1		22-05-2001
<hr/>						
WO 0078731	A	28-12-2000	AU	5548800 A		09-01-2001
			BR	0011770 A		05-03-2002
			CZ	20014583 A3		15-05-2002
			DE	10084704 T0		29-05-2002
			EP	1187816 A1		20-03-2002
			WO	0078731 A1		28-12-2000
			GB	2369360 A		29-05-2002
			NO	20016162 A		18-02-2002
<hr/>						